BIBLIOGRAPHY OF JOURNAL ARTICLES (JANUARY TO JUNE 1998)

PREPARED BY RICHARD R. BROOKMAN, M.D.

CARDIOLOGY

Aldén B, Gilljam T, Gillberg C. Long-term psychological outcome of children after surgery for transposition of the great arteries. Acta Paediatr 1998 Apr; 87(4):405–10. (Study of 31 patients ages 7–21 years.)

Berenson GS, Srinivasan SR, Bao W, et al. Association between multiple cardiovascular risk factors and atherosclerosis in children and young adults. N Engl J Med 1998 Jun; 338(23): 1650–6. (Correlation of autopsy findings and antemortem risk factors in 93 patients ages 2–39 years, most with deaths from trauma.)

Caputo JL, Rudolph DL, Morgan DW. Influence of positive life events on blood pressure in adolescents. J Behav Med 1998 Apr; 21(2):115–29. (Study of 69 sixth-grade students.)

Del Rosario JD, Treiber FA, Harshfield GA, et al. Predictors of future ambulatory blood pressure in youth. J Pediatr 1998 Apr; 132(4):693–8. (Longitudinal study of 88 healthy youth ages 7–13 years followed for 2–3 years.)

Ewart CK, Young DR, Hagberg J. Effects of school-based aerobic exercises on blood pressure in adolescent girls at risk for hypertension. Am J Public Health 1998 Jun; 88(6):949–51. (Study of 88 ninth-grade students.)

Grünberg H, Thetloff M. The cardiovascular risk factor profile of Estonian school children. Acta Paediatr 1998 Jan; 87(1):37–42. (Cross-sectional study of 1018 students ages 9, 12, or 15 years.)

Hulman S, Kushner H, Katz S, et al. Can cardiovascular risk be predicted by newborn, childhood, and adolescent body size? An examination of longitudinal data in urban African Americans. J Pediatr 1998 Jan; 132(1):90–7. (Study of 137 subjects with nine assessments between birth and age 26 to 30 years.)

Katzmarzyk PT, Malina RM, Song TMK, et al. Physique and echocardiographic dimensions in children, adolescents, and young adults. Ann Hum Biol 1998 Mar–Apr; 25(2):145–57. (Study of 143 children ages 8–11 years, 216 youth ages 12–15 years, and 190 young adults ages 16–24 years.)

Lloyd T, Chinchilli VM, Rollings N, et al. Fruit consumption, fitness, and cardiovascular health in female adolescents: The Penn State Young Women's Health Study. Am J Clin Nutr 1998 Apr; 67(4):624–30. (Cross-sectional study of 86 subjects at age 16–18 years.)

Meininger JC, Hayman LL, Coates PM, et al. Genetic and environmental influences on cardiovascular disease risk factors in adolescents. Nurs Res 1998 Jan/Feb; 47(1):11–18. (Comparison of 56 monozygotic and 29 same-sex dizygotic twin pairs at mean age of 12.6 years.)

O'Connor FG, Kugler JP, Oriscello RG. Sudden death in young athletes: Screening for the needle in the haystack. Am Fam Physician 1998 Jun; 57(11):2763–70. (Review of incidence, etiology, assessment of cardiac problems with recommendations of American Heart Association.)

Schmidt GJ, Walkuski JJ, Stensel DJ. The Singapore Youth Coronary Risk and Physical Activity Study. Med Sci Sports Exerc 1998 Jan; 30(1):105–13. (Study of 1579 students ages 6–18 years from six representative districts.)

Stergioula SA, Tripolitsioti A, Messinis D, et al. The effects of endurance training on selected coronary risk factors in children. Acta Paediatr 1998 Apr; 87(4):401–4. (Study of 18 healthy, sedentary Greek boys ages 10–14 years.)

Van Lenthe FJ, van Mechelen W, Kemper HCG. Association of a central pattern of body fat with blood pressure and lipoproteins from adolescence into adulthood. The Amsterdam Growth and Health Study. Am J Epidemiol 1998 Apr; 147(7): 686–93. (Longitudinal study of 84 males, 98 females assessed at mean ages of 13, 14, 15, 16, 21, and 27 years.)

CONTRACEPTION

Brindis C, Boggess J, Katsuranis F, et al. A profile of the adolescent male family planning client. Fam Plann Perspect 1998 Mar/Apr; 30(2):63–6, 88. (Survey of 1540 sexually experienced males ages 19 years and younger.)

Chasan-Taber L, Stampfer MJ. Epidemiology of oral contraceptives and cardiovascular disease. Ann Intern Med 1997 Mar; 128(5):467–77. (Literature review.)

Chiou VM, Shrier LY, Emans SJ. Emergency postcoital contraception. J Pediatr Adolesc Gynecol 1998 May; 11(2):61–72. (Review including discussion of barriers to use, role of health care providers.)

Dias PJ, O'Mara N. Norplant failure: An adolescent case study and review of the literature. J Pediatr Adolesc Gynecol 1998 Feb; 11(1):33–7. (Report of 15-year-old diagnosed with pregnancy 19 months after insertion of Norplant.)

Dull P, Blythe MJ. Preventing teenage pregnancy. Prim Care 1998 Mar; 25(1):111–22. (Review of risk factors, methods of contraception.)

Finer LB, Zabin LS. Does the timing of the first family planning visit still matter? Fam Plann Perspect 1998 Jan/Feb; 30(1):30–3, 42. (Analysis of data from the 1982, 1988, and 1995 cycles of the National Survey of Family Growth for 3252 women younger than 25 years.)

Gold MA, Coupey SM. Young women's attitudes toward injectable and implantable contraceptives. J Pediatr Adolesc Gynecol 1998 Feb; 11(1):17–24. (Survey of 328 patients ages 13–21 years at 3 clinical sites.)

Green TJ, Houghton LA, Donovan U, et al. Oral contraceptives did not affect biochemical folate indexes and homocysteine concentrations in adolescent females. J Am Diet Assoc 1998 Jan; 98(1):49–55. (Study of 229 Ontario women ages 14–20 years.)

O'Dell CM, Forke CM, Polaneczky MM, et al. Depotmedroxyprogesterone acetate or oral contraception in postpartum adolescents. Obstet Gynecol 1998 Apr; 91(4):609–14. (Retrospective study of 161 patients ages 19 years and younger followed at 12 to 18 months postpartum.)

Porter LS. Reducing teenage and unintended pregnancies through client-centered and family-focused school-based family planning programs. J Pediatr Nurs 1998 Jun; 13(3):158–63. (Discussion of theoretical framework and proposed strategy.)

Schuster MA, Bell RM, Berry SH, et al. Impact of a high school condom availability program on sexual attitudes and behaviors. Fam Plann Perspect 1998 Mar/Apr; 30(2):67–72, 88.

© Society for Adolescent Medicine, 2000
Published by Elsevier Science Inc., 655 Avenue of the Americas, New York, NY 10010
1054-139X/00/$–see front matter
PII S1054-139X(00)00107-5

(Anonymous survey of 1945 Los Angeles County students in grades 9–12 with one-year follow-up of 1110 students.)

Thomas BH, DiCenso A, Griffith L. Adolescent sexual behaviour: Results from an Ontario sample. Part II: Adolescent use of protection. Can J Public Health 1998 Mar–Apr; 89(2):94–7. (Longitudinal study of 3272 students ages 10–16 years surveyed annually for five years.)

DERMATOLOGY

DeGroot AC. Fatal attractiveness. The shady side of cosmetics. Clin Dermatol 1998 Jan/Feb; 16(1):167–79. (Review of contact reactions to numerous products.)

Freyre EA, Rebaza RM, Sambi DA, et al. The prevalence of facial acne in Peruvian adolescents and its relation to their ethnicity. J Adolesc Health 1998 Jun; 22(6):480–4. (Cross-sectional study of 2214 healthy youth ages 12–18 years.)

Helm TN, Bergfield WF. Sports dermatology. Clin Dermatol 1998 Jan/Feb; 16(1):159–65. (Illustrated brief review of numerous disorders.)

Rabinowitz LG. Acne vulgaris. AM:STARS 1997 Feb; 8(1):77–86. (Review of basic science, diagnosis, management.)

Strasburger VC. Acne: What every pediatrician should know about treatment. Pediatr Clin North Am 1997 Dec; 44(6):1505–23. (Illustrated review of pathophysiology, management.)

DEVELOPMENTAL DISABILITY

Apajasalo M, Sintonen H, Rautonen J, et al. Health-related quality of life of patients with genetic skeletal dysplasias. Eur J Pediatr 1998 Feb; 157(2):114–21. (Study of 121 adults ages 16–54 years and 19 youth ages 12–15 years.)

Biederman J, Faraone SV, Taylor A, et al. Diagnostic continuity between child and adolescent ADHD: Findings from a longitudinal clinical sample. J Am Acad Child Adolesc Psychiatry 1998 Mar; 37(3):305–13. (Study of 140 affected and 120 control males ages 6–17 years and their families with one- and four-year follow-ups.)

Charney DS. Current issues in attention deficit disorder. J Clin Psychiatry 1998; 59:Suppl 7. (Collected review papers on neurobiology, diagnosis, comorbidities, and psychopharmacologic treatments.)

Goldman LS, Genel M, Bezman RJ, et al. Diagnosis and treatment of attention-deficit/hyperactivity disorder in children and adolescents. JAMA 1998 Apr; 279(14):1100–7. (Council on Scientific Affairs report based on review of literature from 1975 to 1997.)

Kennedy SE, Garcia Martin SD, Kelley JM, et al. Identification of medical and nonmedical needs of adolescents and young adults with spina bifida and their families: A preliminary study. Child Health Care 1998 Winter; 27(1):47–61. (Focus group study of 19 patients ages 12–25 years and needs assessment survey of 54 patients and 77 family members.)

Mannuzza S, Klein RG, Bessler A, et al. Adult psychiatric status of hyperactive boys grown up. Am J Psychiatry 1998 Apr; 155(4): 493–8. (Prospective longitudinal study of 85 probands and 73 controls from ages 5–11 to mean age 25 years.)

Plioplys AV, Kasnicka I, Lewis S, et al. Survival rates among children with severe neurologic disabilities. South Med J 1998 Feb; 91(2):161–72. (Analysis of data collected over 10 years from three skilled nursing facilities for 447 patients ages 1 to 42 years.)

Robin AL. Attention deficit hyperactivity disorder. AM:STARS 1998 Jun; 9(2):373–83. (Review including conceptualization, diagnosis, treatment approaches.)

Saito N, Ebara S, Ohotsuka K, et al. Natural history of scoliosis in spastic cerebral palsy. Lancet 1998 Jun; 351(9117):1687–92. (Study of 37 patients ages 1–15 years followed for up to 10 years.)

Shaywitz SE. Current concepts: Dyslexia. N Engl J Med 1998 Jan; 338(5):307–12. (Review of influencing factors, assessment, management.)

Strauss D, Shavelle R. Life expectancy of adults with cerebral palsy. Dev Med Child Neurol 1998 Jun; 40(6):369–75. (Analysis of vital statistics for 24,768 affected individuals ages 15 years and older from 1980 to 1995.)

ENDOCRINOLOGY

Adams R, Hinkebein MK, McQuillen M, et al. Prompt differentiation of Addison's disease from anorexia nervosa during weight loss and vomiting. South Med J 1998 Feb; 91(2):208–11. (Report of 18-year-old white female admitted to psychiatry service, found to have adrenal insufficiency.)

Åman J, Kroon M, Jones I, et al. Urinary growth hormone excretion in post-menarcheal adolescent girls with type I diabetes. Acta Paediatr 1998 Feb; 87(2):149–53. (Study of 64 diabetic and 64 healthy girls ages 13–18 years.)

Bertelloni S, Paroncelli GI, Sorrentino MC, et al. Effect of central precocious puberty and gonadotropin-releasing hormone analogue treatment on peak bone mass and final height in females. Eur J Pediatr 1998 May; 157(5):363–7. (Study of 14 Italian girls ages 12–14 years.)

Buchlis JG, Irizarry L, Crotzer BC, et al. Comparison of final heights of growth hormone-treated vs untreated children with idiopathic growth failure. J Clin Endocrinol Metab 1998 Apr; 83(4):1075–9. (Study of 94 patients ages 4–17 years.)

Cornean RE, Hindmarsh PC, Brook CGD. Obesity in 21-hydroxylase deficient girls. Arch Dis Child 1998 Mar; 78(3): 261–3. (Retrospective mixed longitudinal study of 22 patients ages 7–20 years.)

Coutant R, Carel J, Letrait M, et al. Short stature associated with intrauterine growth retardation: Final height of untreated and growth hormone-treated children. J Clin Endocrinol Metab 1998 Apr; 83(4):1070–4. (Comparison of 70 treated and 40 untreated French patients at final height at ages 13–19 years.)

Dacou-Voutetakis C, Karavanaki-Karanassiou K, Petrou V, et al. The growth pattern and final height of girls with Turner syndrome with and without human growth hormone treatment. Pediatrics 1998 Apr; 101(4):663–8. (Study of 123 patients.)

Hallidin MU, Tylleskärk K, Hagenäs L, et al. Is growth hormone hypersecretion in diabetic adolescent girls also a daytime problem? Clin Endocrinol 1998 Jun; 48(6):785–94. (Comparison of 11 poorly controlled diabetics and 10 non-diabetic controls ages 13–19 years.)

Harmon RJ, Bender BG, Linden MG, et al. Transition from adolescence to early adulthood: Adaptation and psychiatric status of women with 47,XXX. J Am Acad Child Adolesc Psychiatry 1998 Mar; 37(3):286–91. (Study of 11 affected women and 9 female subling controls at ages 14–19 and at 18–29 years.)

Kordonouri O, Daune T, Enders I, et al. Does the long-term clinical course of type I diabetes mellitus differ in patients with prepubertal and pubertal onset? Results of the Berlin Retinopathy Study. Eur J Pediatr 1998 Mar; 157(3):202–7. (Longitudinal study of 320 patients with onset at ages 1–14 years and 133 patients with onset at ages 9–17 years followed for median of 9 years.)

Leger J, Livoni C, Collin D, et al. Prediction factors in the determination of final height in subjects born small for gestational age. Pediatr Res 1998 Jun; 43(6):808–12. (Longitudinal comparison of 213 SGA and 272 AGA French youth ages 18–22 years.)

Liss DS, Waller DA, Kennard BD, et al. Psychiatric illness and family support in children and adolescents with diabetic ketoacidosis: A controlled study. J Am Acad Child Adolesc

Psychiatry 1998 May; 37(5):536–44. (Comparison of 25 hospitalized and 25 matched ambulatory patients ages 9–17 years.)
Malamitsi-Puchner A, Sarandakou A, Dafogianni C, et al. Serum angiorenin levels in children and adolescents with insulin-dependent diabetes mellitus. Pediatr Res 1998 Jun; 43(6):798–800. (Study of 40 patients and 30 healthy controls ages 5–21 years, Tanner stages 1–5.)
Palardy N, Greening L, Ott J, et al. Adolescents' health attitudes and adherence to treatment for insulin-dependent diabetes mellitus. J Dev Behav Pediatr 1998 Feb; 19(1):31–7. (Study of 101 patients ages 11–18 years attending summer camp.)
Rosenfeld RG, Attie KM, Frane J, et al. Growth hormone therapy of Turner's syndrome: Beneficial effect on adult height. J Pediatr 1998 Feb; 132(2):319–24. (Longitudinal study of 60 patients and 25 controls ages 4–12 years followed to adult height at ages 18–29 years.)
Sandberg DE, MacGillivray MH, Clopper RR, et al. Quality of life among formerly treated childhood-onset growth-hormone deficient adults: A comparison with unaffected siblings. J Clin Endocrinol Metab 1998 Apr; 83(4):1134–42. (Comparison of 140 patients and 53 same-sex siblings ages 18–47 years.)
Weimann E, Bergmann S, Böhles HJ. Oestrogen treatment of constitutional tall stature: A risk-benefit ratio. Arch Dis Child 1998 Feb; 78(2):148–51. (Report of 50 girls, mean age 12 years, treated for up to 2 years.)
Young RS, Rosenbloom AL. N.I.D.D.M. in Minority Children. Clin Pediatr 1998 Feb; 37(2):63–152. (Collected papers from 1996 conference discussing diagnostic criteria, various populations, treatment, prevention, research priorities.)

GYNECOLOGY

Chau JPC, Chang AM, Chang AMZ. Relationship between premenstrual tension syndrome and anxiety in Chinese adolescents. J Adolesc Health 1998 Mar; 22(3):247–9. (Survey of 153 high school students ages 14–18 years.)
Cleckner-Smith CS, Doughty AS, Grossman JA. Premenstrual symptoms: Prevalence and severity in an adolescent sample. J Adolesc Health 1998 May; 22(5):403–8. (Survey of 75 students ages 13–18 years.)
Dealy MF. Dysfunctional uterine bleeding in adolescents. Nurse Pract 1998 May; 23(5):12–23. (Review of diagnosis, clinical assessment, etiologies, treatment.)
Gell JS, Bradshaw KD. Recognition and management of congenital reproductive anomalies. Curr Probl Obstet Gynecol Fertil 1998 May/Jun; 21(3):65–96. (Review.)
Golomb LM, Solidum AA, Ware MP. Primary dysmenorrhea and physical activity. Med Sci Sports Exerc 1998 Jun; 30(6):906–9. (Meta-analysis of seven studies involving high school students, college students, and student nurses.)
Kanizsai B, Örley J, Szigetvári I, et al. Ovarian cysts in children and adolescents: Their occurrence, behavior, and management. J Pediatr Adolesc Gynecol 1998 May; 11(2):85–8. (Study of 119 Hungarian girls ages 5–18 years.)
Levitt MA, Stein DM, Pena A. Gynecologic concerns in the treatment of teenagers with cloaca. J Pediatr Surg 1998 Feb; 33(2):188–93. (Report of 22 patients ages 14 years and older.)
Lubianca JN, Gordon CM, Laufer MR. "Add-back" therapy for endometriosis in adolescents. J Reprod Med 1998 Mar; 43(3): 164–72. (Review with emphasis on GnRH-agonists and bone mass.)
Nikolaev VV, Bizhanova DA. Perineal reconstruction in girls with high vaginal atresia. J Urol 1998 Jun; 159(6):2140–2. (Report of five Russian patients ages 12–14 years with secondary hematocolpos.)
Preminger MK, Pokorny SF. Vaginal discharge—a common pediatric complaint. Contemp Pediatr 1998 Apr; 15(4):115–22. (Review.)
Smith YR, Quint EH, Heitzberg RB. Menorrhagia in adolescents requiring hospitalization. J Pediatr Adolesc Gynecol 1998 Feb; 11(1):13–5. (Retrospective chart review of 37 patients ages 10–20 years.)
Spence JEH. Vaginal and uterine anomalies in the pediatric and adolescent patient. J Pediatr Adolesc Gynecol 1998 Feb; 11(1): 3–11. (Illustrated review.)

HEALTH CARE DELIVERY

Blum RW. Healthy youth development as a model for youth health promotion. A review. J Adolesc Health 1998 May; 22(5):368–75. (Review focusing on risk, stress, and resilience.)
Boekeloo BO, Schamus LA, Simmens SJ, et al. Ability to measure sensitive adolescent behavior via telephone. Am J Prev Med 1998 Apr; 14(3):209–16. (Survey and three month follow-up of 207 youth ages 12–15 years.)
Brener ND, Collins JL. Co-occurrence of health-risk behaviors among adolescents in the United States. J Adolesc Health 1998 Mar; 22(3):209–13. (Analysis of data from 1992 Youth Risk Behavior Survey for 10,645 students ages 12–21 years.)
Ehrman WG, Matson SC. Approach to assessing adolescents on serious or sensitive issues. Pediatr Clin North Am 1998 Feb; 45(1):189–204. (Review of developmental, confidentiality, and consent issues with suggested interviewing techniques.)
Ellen JM, Franzgrote M, Irwin CE, et al. Primary care physicians' screening of adolescent patients: A survey of California physicians. J Adolesc Health 1998 Jun; 22(6):433–8. (Survey of 343 pediatricians, internists, and family physicians concerning care of patients ages 11–14 and 15–18 years.)
Elster AB. Comparison of recommendations for adolescent clinical preventive services developed by national organizations. Arch Pediatr Adolesc Med 1998 Feb; 152(2):193–8. (Comparison of five sets of recommendations for immunizations, health guidance, screening/counseling, and laboratory assessments.)
Ensign J, Santelli J. Health status and service use: Comparison of adolescents at a school-based health clinic with homeless adolescents. Arch Pediatr Adolesc Med 1998 Jan; 152(1):20–4. (Study of 109 youth in emergency shelters and 1010 students using school-based clinics, ages 12–17 years.)
Feinstein RA, Lampkin A, Lorish CD, et al. Medical status of adolescents at time of admission to a juvenile detention center. J Adolesc Health 1998 Mar; 22(3):190–6. (Study of 819 detainees ages 9–19 years.)
Freed LH, Ellen JM, Irwin CE, et al. Determinants of adolescents' satisfaction with health care providers and intentions to keep follow-up appointments. J Adolesc Health 1998 Jun; 22(6): 475–9. (Survey of 124 patients ages 12–21 years before and after a health care visit.)
Grace TW. Health problems of late adolescence. Prim Care 1998 Mar; 25(1):237–52. (Review of screening, health-risk behaviors, common medical problems.)
Gusella JL, Ward A, Butler GS. The experience of hospitalized adolescents: How well do we meet their developmental needs? Child Health Care 1998 Spring; 27(2):131–45. (Survey of 69 Nova Scotian hospitalized patients ages 12–18 years.)
Hagquist CEI. Economic stress and perceived health among adolescents in Sweden. J Adolesc Health 1998 Mar; 22(3): 250–7. (Survey of 2426 ninth-grade students.)
Jepson L, Juszczak L, Fisher M. Mental health care in a high school based health service. Adolescence 1998 Spring; 33(129):1–15. (Description of service and data from 1002 visits by students in grades 9–12.)
Michaud PA, Blum RW, Ferron C. "Bet you I will!" Risk or experimental behavior during adolescence? Arch Pediatr Adolesc Med 1998 Mar; 152(3):224–6. (Commentary.)
Michaud PA, Delbos-Piot I, Narring F. Silent dropouts in health surveys: Are nonrespondent absent teenagers different from

those who participate in school-based health surveys? J Adolesc Health 1998 Apr; 22(4):326–33. (Comparison of 96 absent and 3324 participant students in the Swiss Multicenter Adolescent Survey on Health.)

Montalto NJ. Implementing the guidelines for adolescent preventive services. Am Fam Physician 1998 May; 57(9):2181–8. (Review of content, theory, and application of AMA Guidelines for Adolescent Preventive Services.)

Niskar AS, Kieszak SM, Holmes A, et al. Prevalence of hearing loss among children 6 to 19 years of age: The Third National Health and Nutrition Examination Survey. JAMA 1998 Apr; 279(14):1071–5. (Analysis of audiometric data from 6166 youth.)

Pastore DR, Diaz A. Cultural and medical issues of Latino adolescents. AM:STARS 1998 Jun; 9(2):315–22. (Review of comprehensive assessment, cultural aspects of treatment.)

Potthoff SJ, Bearinger LH, Skay CL, et al. Dimensions of risk behaviors among American Indian youth. Arch Pediatr Adolesc Med 1998 Feb; 152(2):157–63. (Analysis of data from National Indian Adolescent Health Survey for 7687 reservation-based American Indian or Alaska Native students in grades 7–12.)

Reif CJ, Elster AB. Adolescent preventive services. Prim Care 1998 Mar; 25(1):1–21. (Review of adolescent behavior, comprehensive assessment, resources.)

Smith BA, Price JO, Nicholson T, et al. Health knowledge of predominantly Mexican-American high school students. J Health Educ 1998 Jan/Feb; 29(1):21–5. (Survey of 237 students.)

Sonis J. Association between duration of residence and access to ambulatory care among Caribbean immigrant adolescents. Am J Public Health 1998 Jun; 88(6):964–6. (Cross-sectional survey of 1370 New York City high school students ages 13–21 years.)

Stein KF, Roeser R, Markus HR. Self-schemas and possible selves as predictors and outcomes of risky behaviors in adolescents. Nurs Res 1998 Mar/Apr; 47(2):96–106. (Survey of 160 8th-grade students with follow-up of 137 in 9th grade.)

Ziv A, Boulet JR, Slap GB. Emergency department utilization by adolescents in the United States. Pediatrics 1998 Jun; 101(6): 987–94. (Analysis of data from 1994 National Hospital Ambulatory Medical Care Survey for 4308 youth ages 11–14 years, 4096 youth ages 15–17 years, and 6372 youth ages 18–21 years.)

Managed Care

English A, Kapphahn C, Perkins J, et al. Meeting the health needs of adolescents in managed care: A background paper. J Adolesc Health 1998 Apr; 22(4):278–92. (Review of effects of managed care, special needs of adolescents, costs and benefits, delivery systems.)

Fine J. Adolescent pregnancy prevention in managed care. Women's Health Issues 1998 May/Jun; 8(3):148–58. (Review of health insurance status, demographics.)

Kaplan DW, Calonge BN, Guernsey BP, et al. Managed care and school-based health centers: Use of health services. Arch Pediatr Adolesc Med 1998 Jan; 152(1):25–33. (Comparison of 240 youth with and 116 youth without access to school-based health care, all members of Kaiser Permanente managed care.)

Robertson LM, Middleman AB. Knowledge of health insurance coverage by adolescents and young adults attending a hospital-based clinic. J Adolesc Health 1998 Jun; 22(6):439–45. (Survey of 830 patients ages 11–24 years.)

Ryan SA, Millstein SG, Kang M, et al. Adolescents' knowledge of their health insurance coverage. J Adolesc Health 1998 Apr; 22(4):293–9. (Survey of 123 urban youth and 93 adolescent-parent pairs.)

Society for Adolescent Medicine. Meeting the health care needs of adolescents in managed care. J Adolesc Health 1998 Apr; 22(4):271–7. (Position paper focusing on access, comprehensive services, financing of care.)

Thrall JS, McCloskey L, Spivak H, et al. Performance of Massachusetts HMOs in providing Pap smear and sexually transmitted disease screening to adolescent females. J Adolesc Health 1998 Mar; 22(3):184–9. (Review of medical records from seven HMOs for 34,415 females ages 15–21 years.)

HEMATOLOGY/ONCOLOGY

Black P, Gutjahr P, Stopfkuchen H. Physical performance in long-term survivors of acute leukaemia in childhood. Eur J Pediatr 1998 Jun; 157(6):464–7. (Study of 56 German patients ages 9–28 years.)

Blacklay A, Eiser C, Ellis A. Development and evaluation of an information booklet for adult survivors of cancer in childhood. Arch Dis Child 1998 Apr; 78(4):340–4. (Needs assessment of 50 survivors ages 14–32 years and description of resulting educational strategy.)

Kopel SJ, Eiser C, Cool P, et al. Brief report: Assessment of body image in survivors of childhood cancer. J Pediatr Psychol 1998 Apr; 23(2):141–7. (Study of 67 patients ages 12–28 years surveyed 3 to 19 years after diagnosis.)

Manne S, Miller D. Social support, social conflict, and adjustment among adolescents with cancer. J Pediatr Psychol 1998 Apr; 23(2):121–30. (Survey of 50 patients ages 12–20 years.)

Parsons JT, Butler R, Kocik S, et al. The role of the family system in HIV risk reduction: Youths with hemophilia and HIV infection and their parents. J Pediatr Psychol 1998 Feb; 23(1): 57–65. (Multistate study of 125 male youths ages 12–25 years and their parents.)

Puukko L-RM, Sammallahti P, Hovi L, et al. Young female survivors of childhood leukaemia do not have increased somatic concerns. Acta Paediatr 1998 Feb; 87(2):169–74. (Comparison of 42 survivors and 69 healthy controls ages 14–22 years.)

Trentadue NO, Kachoyeanos MK, Lea G. A comparison of two regimens of patient-controlled analgesia for children with sickle cell disease. J Pediatr Nurs 1998 Feb; 13(1):15–9. (Record review of 60 admissions for 26 patients ages 11–18 years.)

Vannatta K, Zeller M, Noll RB, et al. Social functioning of children surviving bone marrow transplantation. J Pediatr Psychol 1998 Jun; 23(3):169–78. (Study of 48 patients and 48 nonchronically ill matched peers ages 8–16 years.)

INFECTIOUS DISEASE

Alderman EM, Shapiro A, Spigland I, et al. Are there risk factors for hepatitis B infection in inner-city adolescents that justify prevaccination screening? J Adolesc Health 1998 May; 22(5): 389–93. (Study of 100 patients attending an urban clinic and 48 patients attending an urban drop-in center, all ages 13–21 years.)

American Academy of Pediatrics Committee on Pediatric AIDS. Human immunodeficiency virus/acquired immunodeficiency syndrome education in schools. Pediatrics 1998 May; 101(5): 933–5. (Position statement focusing on format, curriculum.)

Henry-Reid LM, Rodriguez F, Bell MA, et al. Youth counseled for HIV testing at school- and hospital-based clinics. J Nat Med Assoc 1998 May; 90(5):287–92. (Study of 1652 inner city youths ages 13–19 years.)

Hopkins GL, Hopp JW, Marshak HPH, et al. AIDS risk among students attending Seventh-day Adventist schools in North America. J School Health 1998 Apr; 68(4):141–5. (Survey of 1748 high school students from 69 parochial schools.)

Jemmott JB, Jemmott LS, Fong GT. Abstinence and safer sex HIV risk-reduction interventions for African American adolescents:

A randomized controlled trial. JAMA 1998 May; 279(19):1529–36. (Controlled study of 659 inner-city youth in grades 6–7 with 3, 6, and 12 month follow-up.)

Kokotailo PK, Koscik RE, Henry BC, et al. Health risk taking and human immunodeficiency virus risk in collegiate female athletes. J Am College Health 1998 May; 46(6):263–8. (Survey of 109 athletes and 462 nonathletes ages 18–25 years.)

Liau A, Zimet GD, Fortenberry JD. Attitudes about human immunodeficiency virus immunization. The influence of health beliefs and vaccine characteristics. Sex Transm Dis 1998 Feb; 25(2):76–81. (Survey of 222 undergraduate college students.)

Mikl J, Sudar Z, Smith PF, et al. HIV infection and high risk behaviour among patients attending an STD referral clinic in Prague, Czech Republic. Sex Transm Inf 1998 Apr; 74(2):128–30. (Comparison of 304 patients ages 11–19 years and 1090 patients ages 20 years and older.)

O'Hara P, Parris D, Fichtner RR, et al. Influence of alcohol and drug use on AIDS risk behavior among youth in dropout prevention. J Drug Educ 1998; 28(2):159–68. (Study of 212 students in grades 9–12.)

Rogers AS, Futterman DK, Moscicki AB, et al. The REACH Project of the Adolescent Medicine HIV/AIDS Research Network: Design, methods, and selected characteristics of participants. J Adolesc Health 1998 Apr; 22(4):300–11. (Project description.)

Ruiz J, Facer M, Sun RK. Risk factors for human immunodeficiency virus infection and unprotected anal intercourse among young men who have sex with men. Sex Transm Dis 1998 Feb; 25(2):100–7. (Study of 836 gay or bisexual California males ages 17–25 years.)

Venier JL, Ross MW, Akande A. HIV/AIDS-related social anxieties in adolescents in three African countries. Soc Sci Med 1998 Feb; 46(3):313–20. (Study of 387 Nigerian, 274 Kenyan, and 313 Zimbabwean students in grade 10.)

NUTRITION

Andersen RE, Crespo CJ, Bartlett SJ, et al. Relationship of physical activity and television watching with body weight and level of fatness among children: Results from the Third National Health and Nutrition Examination Survey. JAMA 1998 Mar; 279(12):938–42. (Analysis of data for 4063 youth ages 8–16 years.)

Attia N, Tamborlane WV, Heptulla R, et al. The metabolic syndrome and insulin-like growth factor-I regulation in adolescent obesity. J Clin Endocrinol Metab 1998 May; 83(5):1467–71. (Comparison of 19 obese and 20 lean adolescents ages 12–17 years and 10 lean adults ages 19–25 years.)

Bar-or O, Foeyt J, Bouchard C, et al. Physical activity, genetic, and nutritional considerations in childhood weight management. Med Sci Sports Exerc 1998 Jan; 30(1):2–10. (Summary of roundtable sponsored by American College of Sports Medicine.)

Guezennec CY, Chalabi H, Bernard J, et al. Is there a relationship between physical activity and dietary calcium intake? A survey in 10,373 young French subjects. Med Sci Sports Exerc 1998 May; 30(5):732–9. (Study of 4577 students, 3148 military personnel, and 2648 athletes ages 7–55 years.)

Harel Z, Riggs S, Vaz R, et al. Adolescents and calcium: What they do and do not know and how much they consume. J Adolesc Health 1998 Mar; 22(3):225–8. (Survey of 1117 ninth grade students.)

Hill JO, Trowbridge FL. *The Causes and Health Consequences of Obesity in Children and Adolescents*. Pediatrics 1998 Mar; 101(3)Suppl 497–574. (Series of papers commissioned by the International Life Sciences Institute including epidemiology, assessment, eating behaviors, physical activity, treatment, research priorities.)

Middleman AB, Vazquez I, Durant RH. Eating patterns, physical activity, and attempts to change weight among adolescents. J Adolesc Health 1998 Jan; 22(1):37–42. (Analysis of data from Massachusetts Youth Risk Behavior Survey for 3055 students in grades 9–12.)

Miller EC, Maropis CG. Nutrition and diet-related problems. Prim Care 1998 Mar; 25(1):193–210. (Review of normal nutrition, dietary habits, obesity, sports nutrition.)

Molnar D, Schutz Y. Fat oxidation in nonobese and obese adolescents: Effect of body composition and pubertal development. J Pediatr 1998 Jan; 132(1):98–104. (Study of 235 control and 159 obese Hungarian youth ages 9–16 years.)

Pietrobelli A, Faith MS, Allison DB, et al. Body mass index as a measure of adiposity among children and adolescents: A validation study. J Pediatr 1998 Feb; 132(2):204–10. (Study of 198 healthy Italian youth ages 5–19 years.)

Rössner S. Childhood obesity and adulthood consequences. Acta Paediatr 1998 Jan; 87(1):1–5. (Review of risks, treatment, future strategies.)

Rosner B, Prineas R, Loggie J, et al. Percentiles for body mass index in U.S. children 5 to 17 years of age. J Pediatr 1998 Feb; 132(2):211–22. (Analysis of data from nine large epidemiologic studies including 66,772 subjects.)

Siega-Riz AM, Carson T, Popkin B. Three squares or mostly snacks—What do teens really eat? A sociodemographic study of meal patterns. J Adolesc Health 1998 Jan; 22(1):29–36. (Analysis of data from Continuing Survey of Food Intake by Individuals, 1989–1991 for 1310 youth ages 11–18 years.)

Story M, Neumark-Sztainer D. Diet and adolescent behavior: Is there a relationship? AM:STARS 1998 Jun; 9(2):283–98. (Review focusing on controversies regarding various nutrients, effects of fasting, diet-behavior research.)

Travers SH, Labarta JI, Gargosky SE, et al. Insulin-like growth factor binding protein-1 levels are strongly associated with insulin sensitivity and obesity in early pubertal children. J Clin Endocrinol Metab 1998 Jun; 83(6):1935–9. (Study of 104 healthy pubertal children ages 9–14 years.)

Van Lenthe FJ, van Mechelen W, Kemper HCG, et al. Behavioral variables and development of a central pattern of body fat from adolescence into adulthood in normal-weight whites: The Amsterdam Growth and Health Study. Am J Clin Nutr 1998 May; 67(5):846–52. (Longitudinal study of 84 males, 98 females assessed at mean ages of 13, 14, 15, 16, 21, and 27 years.)

Willi SM, Oexmann MJ, Wright NM, et al. The effects of a high-protein, low-fat, ketogenic diet on adolescents with morbid obesity: Body composition, blood chemistries, and sleep abnormalities. Pediatrics 1998 Jan; 101(1):61–7. (Study of six patients ages 12–15 years weighing 120–198 Kg.)

Eating Disorders

Geist R, Davis R, Heinmaa M. Binge/purge symptoms and comorbidity in adolescents with eating disorders. Can J Psychiatry 1998 Jun; 43(5):507–12. (Study of 120 consecutive patients ages 12–17 years.)

Gila A, Castro J, Toro J, et al. Subjective body-image dimensions in normal and anorexic adolescents. Br J Med Psychol 1998 Jun; 71(2):175–84. (Comparison of 85 anorexic and 427 healthy Spanish females ages 12–18 years.)

Kohn MR, Golden NH, Shenker IR. Cardiac arrest and delirium: Presentations of the refeeding syndrome in severely malnourished adolescents with anorexia nervosa. J Adolesc Health 1998 Mar; 22(3):239–43. (Report of three patients ages 12, 13, and 19 years.)

Neumark-Sztainer D, Story M, Dixon LB, et al. Adolescents engaging in unhealthy weight control behaviors: Are they at risk for other health-compromising behaviors? Am J Public Health 1998 Jun; 88(6):952–5. (Analysis of data from 1993 Youth Risk Behavior Survey for 16,296 students in grades 9–12.)

Neumark-Sztainer D, Story M. Dieting and binge eating among adolescents: What do they really mean? J Am Diet Assoc 1998 Apr; 98(4):446–50. (Focus group study of 138 female, 65 male students in grades 7–12.)

Schwitzer AM, Borgholz K, Dore T, et al. Eating disorders among college women: Prevention, education, and treatment responses. J Am College Health 1998 Mar; 46(5):199–207. (Review of diagnostic criteria, common characteristics, developmental aspects.)

Steiner H, Lock J. Anorexia nervosa and bulimia nervosa in children and adolescents: A review of the past 10 years. J Am Acad Child Adolesc Psychiatry 1998 Apr; 37(4):352–9. (Review of epidemiology, comorbidities, risk factors, prognosis, treatment.)

Wiseman CV, Harris WA, Halmi KA. Eating disorders. Med Clin North Am 1998 Jan;82(1):145–59. (Review of medical consequences, etiology, treatment.)

PREGNANCY/PARENTHOOD

Brabin L, Verhoeff FH, Kazembe P, et al. Improving antenatal care for pregnant adolescents in Southern Malawi. Acta Obstet Gynecol Scand 1998 Apr; 77(4):402–9. (Study of 615 patients ages 10–19 years.)

American Academy of Pediatrics Committee on Adolescence. Counseling the adolescent about pregnancy options. Pediatrics 1998 May; 101(5):938–40. (Position statement on identification, communication, management.)

Deal LW, Holt VL. Young maternal age and depressive symptoms: Results from the 1988 National Maternal and Infant Health Survey. Am J Public Health 1998 Feb; 88(2):266–70. (Comparison of 447 mothers ages 15–17 years, 479 mothers ages 18–19 years, and 870 mothers ages 25–34 years.)

Field TM, Scafidi F, Pickens J, et al. Polydrug-using adolescent mothers and their infants receiving early intervention. Adolescence 1998 Spring; 33(129):117–43. (Study of 126 mothers ages 16–21 years.)

Fiscella K, Kitzman HJ, Cole RE, et al. Does child abuse predict adolescent pregnancy? Pediatrics 1998; 101(4):620–4. (Study of 1026 primiparous African American women ages 17–18 years.)

Gessner BD, Perham-Hester KA. Experience of violence among teenage mothers in Alaska. J Adolesc Health 1998; 22(5):383–8. (Analysis of data from Alaska Pregnancy Risk Assessment Monitoring System for 1991–1994 comparing 910 teenages mothers and 6268 mothers ages 20 years and older.)

Gould J, Blackwell T, Heilig C, et al. Utility of percentage of births to teenagers as a surrogate for the teen birth rate. Am J Public Health 1998 Jun; 88(6):908–12. (Comparison of reporting methods for 1346 mothers ages 10–14 years, 23,983 mothers ages 15–17 years, and 43,073 mothers ages 18–19 years.)

Hellerstedt WL, Story M. Adolescent satisfaction with postpartum contraception and body weight concerns. J Adolesc Health 1998 Jun; 22(6):446–52. (Focus group study of 22 young mothers ages 13–19 years.)

Koshar JH, Lee KA, Goss G, et al. The Hispanic teen mother's origin of birth, use of prenatal care, and maternal and neonatal complications. J Pediatr Nurs 1998 Jun; 13(3):151–7. (Study of 129 pregnant Mexican-Americans ages 14–19 years.)

Lemen PM, Wigten TR, Miller-McCarthy AJ, et al. Screening for gestational diabetes mellitus in adolescent pregnancies. Am J Obstet Gynecol 1998 Jun; 178(6):1251–6. (Retrospective review of 509 patients mean age 16 years.)

Manlove J. The influence of high school dropout and school disengagement on the risk of school-age pregnancy. J Res Adolesc 1998; 8(2):187–220. (Analysis of data from National Educational Longitudinal Study of 1988 for 822 teenaged mothers.)

Morrison DM, Spencer MS, Gillmore MR. Beliefs about substance use among pregnant and parenting adolescents. J Res Adolesc 1998; 8(1):69–95. (Study of 241 girls ages 12–17 years during pregnancy and at 6 and 12 months postpartum.)

Rickert VI, Wiemann CM, Goodrum LA, et al. Employment and health-risk behaviors among pregnant adolescents. J Pediatr Adolesc Gynecol 1998 May; 11(2):79–84. (Multidisciplinary study of 384 girls ages 12–17 years.)

Rome ES, Rybicki LA, Durant RH. Pregnancy and other risk behaviors among adolescent girls in Ohio. J Adolesc Health 1998 Jan; 22(1):50–5. (Analysis of data from 1993 Ohio Youth Risk Behavior Survey for 2314 students in grades 9–12.)

Skinner JD, Pope JF, Carruth B. Alterations in adolescents' sensory taste preferences during and after pregnancy. J Adolesc Health 1998 Jan; 22(1):43–9. (Study of 52 girls ages 11–17 years during third trimester and at 11–13 months postpartum.)

Stevenson W, Maton KI, Teti DM. School importance and dropout among pregnant adolescents. J Adolesc Health 1998 May; 22(5):376–82. (Comparison of 51 Caucasian and 68 African American prenatal patients ages 13–18 years.)

Stouthamer-Loeber M, Wei EH. The precursors of young fatherhood and its effect on delinquency of teenage males. J Adolesc Health 1998 Jan; 22(1):56–65. (Longitudinal study of 506 inner city males followed from age 12–13 years to early 20s.)

Teagle SE, Brindis CD. Substance use among pregnant adolescents: A comparison of self-reported use and provider perception. J Adolesc Health 1998 Mar; 22(3):229–38. (Cross-sectional survey of 248 consecutive prenatal patients ages 15–19 years and interviews with 23 providers.)

Wiemann CM, Berenson AB. Factors associated with recent and discontinued alcohol use by pregnant adolescents. J Adolesc Health 1998 May; 22(5):417–23. (Survey of 378 prenatal patients age 17 years and younger.)

Woods JR. *Substance Abuse in Pregnancy*. Obstet Gynecol Clin North Am 1998 Mar; 25(1):1–272. (Collected reviews on pharmacologic, diagnostic, legal, ethical issues and risks of many specific substances.)

PSYCHIATRY

Bardone AM, Moffitt TE, Caspi A, et al. Adult physical health outcomes of adolescent girls with conduct disorder, depression, and anxiety. J Am Acad Child Adolesc Psychiatry 1998 Jun; 37(6):594–601. (Study of 118 New Zealand girls with a mental health disorder and 341 girls without a disorder at ages 15 and 21 years.)

Berard RMF, Sennett JPB, Ahmed N. Pathways to mental health care among South African adolescents: Analysis of referrals to an adolescent psychiatric unit. Adolescence 1998 Summer; 33(130):405–13. (Study of 670 patients ages 14–25 years.)

Cuffe SP, Addy CL, Garrison CZ, et al. Prevalence of PTSD in a community sample of older adolescents. J Am Acad Child Adolesc Psychiatry 1998 Feb; 37(2):147–54. (Study of 490 youth ages 16–22 years.)

Friedman SB, DeMaso DR. *Adolescent Psychiatric and Behavioral Disorders*. AM:STARS 1998 Jun; 9(2):197–423. (Collected reviews including basic neuroscience, psychopharmacology, problems of displaced youth, depression, conduct disorder, and psychosomatic disorders.)

Jordan KM, Landis DA, Downey MC, et al. Chronic fatigue syndrome in children and adolescents: A review. J Adolesc Health 1998 Jan; 22(1):4–18. (Review of typical clinical features, epidemiology, differential diagnosis, assessment, management.)

Kolter KM, Wilkes TCR, Atkinson M. The prevalence of posttraumatic stress disorder in an adolescent inpatient unit. Can J Psychiatry 1998 Feb; 43(1):64–8. (Two-year retrospective study of 187 patients ages 13–19 years.)

Li X, Howard D, Stanton B, et al. Distress symptoms among urban African American children and adolescents: A psychometric evaluation of the Checklist of Children's Distress Symptoms. Arch Pediatr Adolesc Med 1998 Jun; 152(6):569–77. (Study of 349 low-income youth ages 9–15 years.)

Pine DS, Cohen P, Gurley D, et al. The risk for early-adulthood anxiety and depressive disorders in adolescents with anxiety and depressive disorders. Arch Gen Psychiatry 1998 Jan; 55(1):56–64. (Longitudinal study of 776 selected subjects ages 9–18 years assessed at baseline and after 2 and 9 years.)

Post D, Carr C, Weigand J. Teenagers: Mental health and psychological issues. Prim Care 1998 Mar; 25(1):181–92. (Review of various disorders.)

Reinecke MA, Ryan NE, DuBois DL. Cognitive-behavioral therapy of depression and depressive symptoms during adolescence: A review and meta-analysis. J Am Acad Child Adolesc Psychiatry 1998 Jan; 37(1):26–34. (Analysis of six studies involving 217 subjects.)

Roberts RE, Attkisson CC, Rosenblatt A. Prevalence of psychopathology among children and adolescents. Am J Psychiatry 1998 Jun; 155(6):715–25. (Literature review of 52 studies from past four decades.)

Schultz SC, Findling RL, Wise A, et al. Child and adolescent schizophrenia. Psychiatr Clin North Am 1998 Mar; 21(1):43–56. (Review of epidemiology, diagnosis, neuropsychiatric research, treatment, family issues.)

Sleegers J, Spijker J, van Limbeck J, et al. Mental health problems among homeless adolescents. Acta Psychiatr Scand 1998 Apr; 97(4):253–9. (Review of 18 surveys of homeless youth ages 16–23 years from various sites in Amsterdam.)

Zametkin AJ, Ernst M, Silver R. Laboratory and diagnostic testing in child and adolescent psychiatry: A review of the past 10 years. J Am Acad Child Adolesc Psychiatry 1998 May; 37(5): 464–72. (Review of neuroendocrinologic tests, neurotransmitter measurements, toxicology, genetic studies, pharmacologic monitoring.)

Suicide

Beautrais AL, Joyce PR, Mulder RT. Psychiatric contacts among youths aged 13 through 24 years who have made serious suicide attempts. J Am Acad Child Adolesc Psychiatry 1998 May; 37(5):504–11. (Comparison of 129 suicide attempters and 153 community controls.)

Bell CC, Clark CC. Adolescent suicide. Pediatr Clin North Am 1998 Apr; 45(2):365–80. (Review of epidemiology, risk factors, resiliency factors, prevention, intervention.)

Gould MS, Shaffer D, Fisher P, et al. Separation/divorce and child and adolescent completed suicide. J Am Acad Child Adolesc Psychiatry 1998 Feb; 37(2):155–62. (Case-controlled psychologic autopsy study of 120 consecutive suicides in persons under age 20 years with 147 community controls.)

Groholt B, Ekeberg O, Wichstrom L, et al. Suicide among children and younger and older adolescents in Norway: A comparative study. J Am Acad Child Adolesc Psychiatry 1998 May; 37(5): 473–81. (Retrospective comparison of 14 children ages 8–14 years and 115 youth ages 15–19 years who completed suicide and 889 controls.)

Harrington R, Kerfoot M, Dyer E, et al. Randomized trial of a home-based family intervention for children who have deliberately poisoned themselves. J Am Acad Child Adolesc Psychiatry 1998 May; 37(5):512–8. (Study of 162 patients ages 10–16 years.)

Hultén A, Wasserman D. Lack of continuity—A problem in the care of young suicides. Acta Psychiatr Scand 1998 May; 97(5): 326–33. (Retrospective chart review of 34 Swedish youth ages 15–24 years who completed suicide.)

Johnson BA, Brent DA, Bridge J, et al. The familial aggregation of adolescent suicide attempts. Acta Psychiatr Scand 1998 Jan; 97(1):18–24. (Comparison of relatives of 62 referred attempters and 70 never suicidal controls ages 14–18 years.)

Laurent A, Foussard N, David M, et al. A 5-year follow-up study of suicide attempts among French adolescents. J Adolesc Health 1998 May; 22(5):424–30. (Study of 552 patients and 293 hospitalized nonpsychiatric patients ages 8–17 years.)

McKeown RE, Garrison CZ, Cuffe SP, et al. Incidence and predictors of suicidal behaviors in a longitudinal sample of young adolescents. J Am Acad Child Adolesc Psychiatry 1998 Jun; 37(6):612–9. (Study of 359 secondary school students ages 11–16 years with two-year follow-up.)

Neugebauer R, Reuss ML. Association of maternal, antenatal, and perinatal complications with suicide in adolescence and young adulthood. Acta Psychiatr Scand 1998 Jun; 97(6):412–8. (Case-controlled retrospective study of 189 youths ages 15–22 years completing suicide.)

Pfeffer CR, Normandin L, Kakuma T. Suicidal children grow up: Relations between family psychopathology and adolescents' lifetime suicidal behavior. J Nerv Ment Dis 1998 May; 186(5): 269–75. (Study of 650 first and 1174 second degree biological relatives of 69 children at risk for suicidal behavior and 61 community controls followed from childhood to age 10–21 years.)

Rubenstein JL, Halton A, Kasten L, et al. Suicidal behavior in adolescents: Stress and protection in different family contexts. Am J Orthopsychiatry 1998 Apr; 68(2):274–84. (Survey of 272 public high school students in grades 10–11.)

Stein D, Apter A, Ratzoni G, et al. Association between multiple suicide attempts and negative affects in adolescents. J Am Acad Child Adolesc Psychiatry 1998 May; 37(5):488–94. (Comparison of 32 first attempters, 19 repeated attempters, 109 nonsuicidal psychiatric inpatients, and 55 community controls ages 12–18 years.)

Stein D, Brom D, Elizur A, et al. The association between attitudes toward suicide and suicidal ideation in adolescents. Acta Psychiatr Scand 1998 Mar; 97(3):195–201. (Survey of 525 Israeli military recruits ages 18–19 years.)

Taiminen TJ, Kallio-Soukainen K, Nokso-Koivisto H, et al. Contagion of deliberate self-harm among adolescent inpatients. J Am Acad Child Adolesc Psychiatry 1998 Feb; 37(2):211–7. (Study of 12 female patients ages 12–18 years.)

PSYCHOLOGY/SOCIOLOGY

Adebayo B, Gambling behavior of students in grades seven and eight in Alberta, Canada. J School Health 1998 Jan; 68(1):7–11. (Survey of 745 students ages 12–15 years.)

Borowsky IW, Resnick MD. Environmental stressors and emotional status of adolescents who have been in special education classes. Arch Pediatr Adolesc Med 1998 Apr; 152(4):377–82. (Analysis of data from 1992 Minnesota Student Survey for 121,848 students in grades 6, 9, and 12.)

Britto MT, Garrett JM, Dugliss MAJ, et al. Risky behavior in teens with cystic fibrosis or sickle cell disease: A multicenter study. Pediatrics 1998 Feb; 101(2):250–6. (Study of 292 patients with sickle cell disease and 146 patients with cystic fibrosis ages 12–19 years and matched controls.)

Earl DT, Blackwelder RB. Management of chronic medical conditions in children and adolescents. Prim Care 1998 Mar; 25(1): 253–70. (Review of common management issues, with examples of specific chronic illnesses and disabling conditions.)

Orr DP. Helping adolescents toward adulthood. Contemp Pediatr 1998 May; 15(5):55–74. (Review of physical, emotional, and social development.)

Reiff MI. Adolescent school failure: Failure to thrive in adolescence. Pediatr Rev 1998 Jun; 19(6):199–207. (Review of ante-

cedents, prevention, differential diagnosis, evaluation, and treatment planning.)

Reynolds AJ. Resilience among black urban youth. Prevalence, intervention effects, and mechanisms of influence. Am J Orthopsychiatry 1998 Jan; 68(1):84–100. (Study of 1170 youth age 12 years.)

Violence

American College of Physicians. Firearm injury prevention. Ann Intern Med 1998 Feb; 128(3):236–41. (Position statement.)

Cirillo KJ, Pruitt BE, Colwell B, et al. School violence: Prevalence and intervention strategies for at-risk adolescents. Adolescence 1998 Summer; 33(130):319–30. (Study of 50 Texas students in grades 9–12.)

Dahlberg LL. Youth violence in the United States. Major trends, risk factors, and prevention approaches. Am J Prev Med 1998 May; 14(4):259–72. (Extensive review of homicide victimization and perpetration in past decade.)

Foshee VA, Bauman KE, Arriaga XB, et al. An evaluation of Safe Dates, an adolescent dating violence prevention program. Am J Public Health 1998 Jan; 88(1):45–50. (Controlled study of 1700 rural students in grades 8–9.)

Ginsberg KR, Fein JA, Johnson CD. Violence prevention in the early years. Contemp Pediatr 1998 Apr; 15(4):97–110. (Review of childhood antecedents of adolescent violence.)

Hennes H, Calhoun AD. *Violence Among Children and Adolescents*. Pediatr Clin North Am 1998 Apr; 45(2):269–474. (Collected reviews including statistics, theoretical frameworks, risk and resiliency factors, media influence, consequences, dating violence, child abuse, corporal punishment.)

Kelton GM, Shank JC. Adolescent injury and death. The plagues of accident, self-infliction, and violence. Prim Care 1998 Mar; 25(1):163–79. (Review with emphasis on violence and prevention strategies.)

Kulig J, Valentine J, Griffith J, et al. Predictive model of weapon carrying among urban high school students: Results and validation. J Adolesc Health 1998 Apr; 22(4):312–9. (Survey of 1189 students in two Boston high schools.)

Kuther TL, Fisher CB. Victimization by community violence in young adolescents from a suburban city. J Early Adolesc 1998 Feb; 18(1):53–76. (Study of 123 students in grades 6–8.)

Lowry R, Powell KE, Kann L, et al. Weapon-carrying, physical fighting, and fight-related injury among U.S. adolescents. Am J Prev Med 1998 Feb; 14(2):122–9. (Analysis of data from 1992 Youth Risk Behavior Survey for 10,269 youth ages 12–21 years.)

Rich M, Woods ER, Goodman E, et al. Aggressors or victims: Gender and race in music video violence. Pediatrics 1998 Apr; 101(4):669–74. (Content analysis of 518 music videos televised during four week period on national networks.)

Simon TR, Richardson JL, Dent CW, et al. Prospective psychosocial, interpersonal, and behavioral predictors of handgun carrying among adolescents. Am J Public Health 1998 Jun; 88(6): 960–3. (Survey of 4370 students in 9th grade with followup of 2421 students in 12th grade.)

Song L, Singer MI, Anglin TM. Violence exposure and emotional trauma as contributors to adolescents' violent behaviors. Arch Pediatr Adolesc Med 1998 Jun; 152(6):531–6. (Survey of 3735 students ages 14–19 years in grades 9–12.)

Synovitz LB, Byrne TJ. Antecedents of sexual victimization: Factors discriminating victims from non-victims. J Am College Health 1998 Jan; 46(4):151–8. (Survey of 241 female college students.)

Tschumper A, Narring F, Meier C, et al. Sexual victimization in adolescent girls (age 15–20 years) enrolled in post-mandatory schools or professional training programmes in Switzerland. Acta Paediatr 1998 Feb; 87(2):212–7. (Analysis of data from Swiss Multicenter Adolescent Survey on Health for 3993 girls of whom 711 reported victimization.)

PUBERTY

Armstrong N, Welsman JR, Kirby BJ. Peak oxygen uptake and maturation in 12-year olds. Med Sci Sports Exerc 1998 Jan; 30(1):165–9. (Study of 106 boys and 106 girls Tanner stages 1–4.)

Cheng JCY, Leung SSF, Chiu BSK, et al. Can we predict body height from segmental bone length measurements? A study of 3647 children. J Pediatr Orthop 1998 May/Jun; 18(3):387–93. (Cross-sectional study of Chinese students ages 3–18 years.)

Cronau H, Brown RT. Growth and development: Physical, mental, and social aspects. Prim Care 1998 Mar; 25(1):23–47. (Review of puberty, cognitive development, psychosocial development, problems related to growth and maturation.)

Eliakim A, Brasel JA, Barstow TJ, et al. Peak oxygen uptake, muscle volume, and the growth hormone-insulin-like growth factor-1 axis in adolescent males. Med Sci Sports Exerc 1998 Apr; 30(4):512–7. (Study of 43 healthy students ages 15–17 years, Tanner stages 3–5.)

Galluzzi F, Salti R, Bindi G, et al. Adult height comparison between boys and girls with precocious puberty after long-term gonadotrophin-releasing hormone analogue therapy. Acta Paediatr 1998 May; 87(5):521–7. (Longitudinal study of 33 Italian patients treated at age 5–8 years followed to adult height.)

Mølgaard C, Thomsen BL, Michaelsen KF. Influence of weight, age and puberty on bone size and bone mineral content in healthy children and adolescents. Acta Paediatr 1998 May; 87(5):494–9. (Study of 201 female, 142 male Danish youth ages 5–19 years.)

Roemmich JN, Clark PA, Mai V, et al. Alterations in growth and body composition during puberty. III. Influence of maturation, gender, body composition, fat distribution, aerobic fitness, and energy expenditure on nocturnal growth hormone release. J Clin Endocrinol Metab 1998 May; 83(5):1440–7. (Controlled research center study of 30 prepubertal and 29 pubertal youth.)

Sher ES, Migeon CJ, Berkavitz GD. Evaluation of boys with marked breast development at puberty. Clin Pediatr 1998 Jun; 37(6):367–72. (Report of 60 patients ages 9–19 years seen during 10 year span.)

Warner JT, Cowan FJ, Dunstan FDJ, et al. Measured and predicted bone mineral content in healthy boys and girls aged 16–18 years: Adjustment for body size and puberty. Acta Paediatr 1998 Mar; 87(3):244–9. (Cross-sectional study of 58 Welsh youth.)

PULMONOLOGY

Gibson PG, Shah S, Mamoon HA. Peer-led asthma education for adolescents: Impact evaluation. J Adolesc Health 1998 Jan; 22(1):66–72. (Study of 92 students with asthma in 2 Australian girls' high schools.)

Huet F, Semama D, Maingueneau C, et al. Vitamin A deficiency and nocturnal vision in teenagers with cystic fibrosis. Eur J Pediatr 1997 Dec; 156(12):949–51. (Prospective study of 10 clinically stable patients ages 12–20 years.)

Lai H-C, Losorok MR, Sondel SA, et al. Growth status in children with cystic fibrosis based on the National Cystic Fibrosis Patient Registry data: Evaluation of various criteria used to identify malnutrition. J Pediatr 1998 Mar; 132(3):478–85. (Cross-sectional analysis of data from 1993 for 13,116 patients ages birth to 18 years followed at 114 centers.)

Morison S, Dodge JA, Cole TJ, et al. Height and weight in cystic fibrosis: A cross-sectional study. Arch Dis Child 1997 Dec; 77(6):497–500. (Study of 3056 patients ages birth to 30 years from 31 large United Kingdom clinics.)

Mullins LL, Chaney JM, Pace TM, et al. Illness uncertainty, attributional style, and psychological adjustment in older adolescents and young adults with asthma. J Pediatr Psychol 1997 Dec; 22(6):871–80. (Study of 49 college students ages 17–26 years.)

Pine DS, Coplan JD, Papp LA, et al. Ventilatory physiology of children and adolescents with anxiety disorders. Arch Gen Psychiatry 1998 Feb; 55(2):123–9. (Pulmonary function testing of 18 patients and 15 healthy controls ages 7–16 years.)

SEXUALITY

Abma J, Driscoll A, Moore K. Young women's degree of control over first intercourse: An exploratory analysis. Fam Plann Perspect 1998 Jan/Feb; 30(1):12–8. (Analysis of data from 1995 National Survey of Family Growth for 2043 women ages 15–24 years.)

Carpenter LM. From girls into women: Scripts for sexuality and romance in *Seventeen* magazine, 1974–1994. J Sex Res 1998; 35(2):158–68. (Quantitative and qualitative content analysis of 244 articles from 1974, 1984, and 1994 issues.)

Dickson N, Paul C, Herbison P, et al. First sexual intercourse: Age, coercion, and later regrets reported by a birth cohort. BMJ 1998 Jan; 316(7124):29–33. (Crosssectional study of 477 male and 458 female New Zealanders at ages 18 and 21 years.)

Downey JI, Friedman RC. Female homosexuality: Classical psychoanalytic theory reconsidered. J Am Psychol Assoc 1998 Jun; 46(2):471–506. (Literature review.)

Faulkner AH, Cranston K. Correlates of same-sex sexual behavior in a random sample of Massachusetts high school students. Am J Public Health 1998 Feb; 88(2):262–6. (Analysis of data from 1993 Youth Risk Behavior Survey for 3054 students in grades 9–12.)

Garofalo R, Wolf RC, Kessel S, et al. The association between health risk behaviors and sexual orientation among a school-based sample of adolescents. Pediatrics 1998 May; 101(5):895–902. (Analysis of data from 1995 Massachusetts Youth Risk Behavior Survey for 4159 students in grades 9–12.)

Haffner DW. Facing facts. Sexual health for American adolescents. J Adolesc Health 1998 Jun; 22(6):453–9. (1997 Gallagher lecture.)

Jejeebhoy SJ. Adolescent sexual and reproductive behaior: A review of the evidence from India. Soc Sci Med 1998 May; 46(10):1275–90. (Review of attitudes and behaviors regarding sexuality, fertility and its control, marriage.)

Langille DB, Andreou P, Beazley RP, et al. Sexual health knowledge of students at a high school in Nova Scotia. Can J Public Health 1998 Mar–Apr; 89(2):85–9. (Survey of 640 students in grades 9–12.)

Miller BC, Norton MC, Fan X, et al. Pubertal development, parental communication, and sexual values in relation to adolescent sexual behaviors. J Early Adolesc 1998 Feb; 18(1): 27–52. (Study of 473 Utah families interviewed in 1991, 1992, and 1993 each with a young adolescent.)

Oliver DP, Leeming FC, Dwyer WO. Studying parental involvement in school-based sex education: Lessons learned. Fam Plann Perspect 1998 May/Jun; 30(3):143–7. (Study of 775 Memphis public school students in grades 5–8 and their parents.)

Savin-Williams RC. The disclosure to families of same-sex attractions by lesbian, gay, and bisexual youths. J Res Adolesc 1998; 8(1):49–68. (Literature review.)

Shrier LA, Pierce JD, Emans SJ, et al. Gender differences in risk behaviors associated with forced or pressured sex. Arch Pediatr Adolesc Med 1998 Jan; 152(1):57–63. (Survey of 7884 sexually active students in grades 8–12 in Vermont.)

Sonenstein FL, Ku L, Lindberg L, et al. Changes in sexual behavior and condom use among teenage males: 1988 to 1995. Am J Public Health 1998 Jun; 88(6):956–9. (Analysis of data from 1988 and 1995 National Surveys of Adolescent Males for 1708 and 1874 youth ages 15–19 years.)

Stephenson JM, Oakley A, Charleston S, et al. Behavioral intervention trials for HIV/STD prevention in schools: Are they feasible? Sex Transm Inf 1998 Jun; 74(6):405–8. (Comparison of peer-led and teacher-led sex education in four Greater London secondary schools involving 469 students.)

Thomas BH, DiCenso A, Griffith L. Adolescent sexual behavior: Results from an Ontario sample. Part I: Adolescent sexual activity. Can J Public Health 1998 Mar–Apr; 89(2):90–3. (Survey of 3272 students ages 10–16 years yearly for 5 years.)

Tobias BB, Ricer RE. Counseling adolescents about sexuality. Prim Care 1998 Mar; 25(1):49–70. (Review of psychosexual development, risk factors, sex education, office assessment and guidance.)

Upchurch DM, Levy-Storms L, Sucoff CA, et al. Gender and ethnic differences in the timing of first sexual intercourse. Fam Plann Perspect 1998 May/Jun; 30(3):121–7. (Longitudinal study of 877 Los Angeles County youth ages 12–17 years.)

SEXUALLY TRANSMITTED DISEASE

Baker JG, Rosenthal SL. Psychological aspects of sexually transmitted infection acquisition in adolescent girls: A developmental perspective. J Dev Behav Pediatr 1998 Jun; 19(3):202–8. (Review of biological, psychosexual, cognitive, sociocultural factors.)

Beck-Sague CM, Farshy CE, Jackson TK, et al. Detection of *Chlamydia trachomatis* cervical infection by urine tests among adolescents clinics. J Adolesc Health 1998 Mar; 22(3):197–204. (Comparison of urine and cervical specimens from 415 patients ages 13–20 years.)

Benaim J, Pulaski M, Coupey SM. Adolescent girls and pelvic inflammatory disease: Experience and practices of emergency department pediatricians. Arch Pediatr Adolesc Med 1998 May; 152(5):449–54. (Survey of 54 physicians.)

Bonny AE, Biro FM. Recognizing and treating STDs in adolescent girls. Contemp Pediatr 1998 Mar; 15(3):119–43. (Review.)

Buzi RS, Weinman ML, Smith PB. Ethnic differences in STD rates among female adolescents. Adolescence 1998 Summer; 33(130): 313–8. (Study of 205 family planning clinic clients ages 13–20 years.)

Clark LR, Brasseux C, Richmond D, et al. Effect of HIV counseling and testing on sexually transmitted diseases and condom use in an urban adolescent population. Arch Pediatr Adolesc Med 1998 Mar; 152(3):269–73. (Study of 149 patients ages 15–19 years with two-year follow-up of 126.)

Darville T. Chlamydia. Pediatr Rev 1998 Mar; 19(3):85–91. (Review of pathogenesis, clinical manifestations, diagnosis, treatment, prevention.)

Edwards S, Carne C. Oral sex and the transmission of non-viral STIs. Sex Transm Inf 1998 Apr; 74(2):95–100. (Review.)

Ellen JM, Aral SO, Madger LS. Do differences in sexual behaviors account for the racial/ethnic differences in adolescents' self-reported history of a sexually transmitted disease? Sex Transm Dis 1998 Mar; 25(3):125–9. (Analysis of data from 1992 Youth Risk Behavior Survey for 5189 sexually experienced United States youth ages 14–21 years.)

Gunn RA, Podschun GD, Fitzgerald S, et al. Screening high-risk adolescent males for *Chlamydia trachomatis* infection. Obtaining urine specimens in the field. Sex Transm Dis 1998 Jan; 25(1): 49–52. (Study of 261 San Diego males using outreach workers ages 16–21 years.)

Ho GYF, Bierman R, Beardsley L, et al. Natural history of cervicovaginal papillomavirus infection in young women. N Engl J Med 1998 Feb; 338(7):423–8. (Study of 608 college women ages 17–23 years followed for 3 years.)

Krowchuk DP. Sexually transmitted diseases in adolescents: What's new? South Med J 1998 Feb; 91(2):123–31. (Review of risk factors, epidemiology, diagnosis, treatment.)

Lappa S, Coleman MT, Moscicki AB. Managing sexually transmitted diseases in adolescents. Prim Care 1998 Mar; 25(1):71–110. (Review of clinical evaluation, prevention, common syndromes.)

Moscicki AB, Shiboski S, Broering J, et al. The natural history of human papillomavirus infection as measured by repeated DNA testing in adolescent and young women. J Pediatr 1998 Feb; 132(2):277–84. (Longitudinal study of 618 women ages 13–22 years seen in family planning or university student health clinic.)

Oh MK, Smith KR, O'Cain M, et al. Urine-based screening of adolescents in detention to guide treatment for gonococcal and chlamydial infections: Translating research into intervention. Arch Pediatr Adolesc Med 1998 Jan; 152(1):52–6. (Cross-sectional screening of 217 male and 46 female detainees.)

Ohlemeyer CL, Hornberger LL, Lynch DA, et al. Diagnosis of *Trichomonas vaginalis* in adolescent females: Inpouch TV culture versus wet-mount microscopy. J Adolesc Health 1998 Mar; 22(3):205–8. (Study of 467 patients ages 12–18 years.)

Sung L, MacDonald NE. Gonorrhea: A pediatric perspective. Pediatr Rev 1998 Jan; 19(1):13–6. (Review of epidemiology, clinical manifestations, management.)

Sung L, MacDonald NE. Syphilis: A pediatric perspective. Pediatr Rev 1998 Jan; 19(1):17–22. (Review of epidemiology, clinical manifestations, laboratory diagnosis, management.)

SPORTS MEDICINE

Baumert PW, Henderson JM, Thompson NJ. Health risk behaviors of adolescent participants in organized sports. J Adolesc Health 1998 Jun; 22(6):460–5. (Survey of 6849 students in grades 9–12 comparing athletes and nonathletes.)

Bowers LD. Athletic drug testing. Clin Sports Med 1998 Apr; 17(2):299–318. (Review of substances used, laboratory methods, consequences.)

Cantu RC. *Neurologic Athletic Head and Neck Injuries*. Clin Sports Med 1998 Jan; 17(1):1–210. (Collected reviews including epidemiology, initial assessment, outcome, medicolegal aspects, injuries in selected impact sports.)

Choma CW, Sforzo GA, Keller BA. Impact of rapid weight loss on cognitive function in collegiate wrestlers. Med Sci Sports Exerc 1998 May; 30(5):746–9. (Study of 14 wrestlers and 15 controls ages 18–22 years.)

Garcia AW, Pender NJ, Antonakos CL, et al. Changes in physical activity beliefs and behaviors in boys and girls across the transition to junior high school. J Adolesc Health 1998 May; 22(5):394–402. (Longitudinal study of 132 students in 6th and again in 7th grades.)

Glover DW, Maron BJ. Profile of preparticipation cardiovascular screening for high school athletes. JAMA 1998 Jun; 279(22): 1817–9. (Analysis of guidelines and requirements from all high school jurisdictions in the United States compared with 1996 American Heart Association consensus guidelines.)

Hergenroeder AC. Prevention of sports injuries. Pediatrics 1998 Jun; 101(6):1057–63. (Review of epidemiology, risk reduction strategies, rehabilitation of injuries.)

Kaeding CC, Whitehead R. Musculoskeletal injuries in adolescents. Prim Care 1998 Mar; 25(1):211–23. (Illustrated review of numerous injuries and conditions.)

Katzmarzyk PT, Malina RM, Song TMK, et al. Physical activity and health-related fitness in youth: A multivariate analysis. Med Sci Sports Exerc 1998 May; 30(5):709–14. (Study of 630 Quebec youth ages 9–18 years.)

Lyon RM, Street CC. Pediatric sports injuries: When to refer or X-ray. Pediatr Clin North Am 1998 Feb; 45(1):221–44. (Review organized by anatomic location.)

Madsen KL, Adams WC, Van Loan MD. Effects of physical activity, body weight and composition, and muscular strength on bone density in young women. Med Sci Sports Exerc 1998 Jan; 30(1):114–20. (Study of 60 eumenorrheic college students ages 18–26 years comparing athletes and nonathletes.)

Myers A, Sickles T. Preparticipation sports examination. Prim Care 1998 Mar; 25(1):225–36. (Review of history, physical examination, use of sample forms.)

Nigg CR, Courneya KS. Transtheoretical model: Examining adolescent exercise behavior. J Adolesc Health 1998 Mar; 22(3): 214–24. (Survey of 819 students in grades 9–12.)

Nindl BC, Kraemer WJ, Emmert WH, et al. Comparison of body composition assessment among lean black and white male collegiate athletes. Med Sci Sports Exerc 1998 May; 30(5):769–76. (Study of 34 white, 30 black athletes ages 11–22 years.)

Page RM, Hammermeister J, Scanlan A, et al. Is school sports participation a protective factor against adolescent health risk behaviors? J Health Educ 1998 May/Jun; 29(3):186–92. (Analysis of data from 1991 Youth Risk Behavior Survey for 12,272 students in grades 9–12.)

Rowland TW. The biological basis of physical activity. Med Sci Sports Exerc 1998 Mar; 30(3):392–9. (Review of evidence for central control of habitual physical activity with implications for preventive health.)

Sturmi JE, Diorio DJ. Anabolic agents. Clin Sports Med 1998 Apr; 17(2):261–82. (Review of pharmacology, clinical uses, athletic use, side effects, legal issues.)

Wroble RR, Moxley DP. Acute weight gain and its relationship to success in high school wrestlers. Med Sci Sports Exerc 1998 Jun; 30(6):949–54. (Study of 260 competitors in a high school wrestling tournament.)

Wroble RR, Moxley DP. Weight loss patterns and success rates in high school wrestlers. Med Sci Sports Exerc 1998 Apr; 30(4): 625–8. (Study of 465 wrestlers from weight classes 103 to 189 pounds.)

SUBSTANCE USE/ABUSE

American Academy of Child and Adolescent Psychiatry. Summary of the practice parameters for the assessment and treatment of children and adolescents with substance use disorders. J Am Acad Child Adolesc Psychiatry 1998 Jan; 37(1):122–6. (Guidelines.)

American Academy of Pediatrics Committee on Substance Abuse. Tobacco, alcohol, and other drugs: The role of the pediatrician in prevention and management of substance abuse. Pediatrics 1998 Jan; 101(1):125–8. (Position statement focusing on pervasiveness of drug use, evaluation, drug testing, management issues.)

Bhattacharya G. Drug use among Asian-Indian adolescents: Identifying protective/risk factors. Adolescence 1998 Spring; 33(129):169–84. (Literature review.)

Black DR, Tobler NS, Sciacca JP. Peer helping/involvement: An efficacious way to meet the challenge of reducing alcohol, tobacco, and other drug use among youth? J School Health 1998 Mar; 68(3):87–93. (Meta-analysis of 120 studies of school-based prevention programs, deriving features of promising model programs.)

Brasseux C, D'Angelo LJ, Guagliardo M, et al. The changing pattern of substance abuse in urban adolescents. Arch Pediatr Adolesc Med 1998 Mar; 152(3):234–7. (Results of blinded, anonymous urine sampling of 1313 patients from 1995 to 1996 and 1312 patients from 1989 to 1990 ages 12–21 years.)

Brooks AJ, Stuewig J, LeCroy CW. A family based model of Hispanic adolescent substance use. J Drug Educ 1998; 28(1):

65–86. (Study of 413 Mexican-American middle school students ages 11–15 years.)

Chou C, Montgomery S, Pentz M, et al. Effects of a community-based prevention program on decreasing drug use in high-risk adolescents. Am J Public Health 1998 Jun; 88(6):944–8. (Comparison of 1904 experimental and 1508 control students on entry to grade 6 or 7 and yearly after 6, 18, 30, and 42 months.)

Crowley TJ, Mikulich SK, MacDonald M, et al. Substance-dependent, conduct-disordered adolescent males: Severity of diagnosis predicts 2-year outcome. Drug Alc Depend 1998 Feb; 49(3):225–37. (Study of 87 youth ages 13–19 years in residential treatment.)

Deas-Nesmith D, Campbell S, Brady KT. Substance use disorders in an adolescent inpatient psychiatric population. J Nat Med Assoc 1998 Apr; 90(4):233–8. (Study of 100 consecutive patients ages 13–17 years.)

Donnermeyer JF, Davis RR. Cumulative effects of prevention education on substance use among 11th grade students in Ohio. J School Health 1998 Apr; 68(4):151–8. (Statewide survey of 3190 students compared by number of exposures to school-based prevention activities.)

Gabel S, Stallings MC, Young SE, et al. Family variables in substance-misusing male adolescents: The importance of maternal disorder. Am J Drug Alc Abuse 1998; 24(1):61–84. (Study of families of 50 patients in residential treatment and 50 controls ages 13–19 years.)

Harrison PA, Fulkerson JA, Beebe TJ. DSM-IV substance use disorder criteria for adolescents: A critical examination based on a statewide school survey. Am J Psychiatry 1998 Apr; 155(4):486–92. (Analysis of data from 1995 Minnesota Student Survey for 74,008 students in grades 9–12.)

Iven VG. Recreational drugs. Clin Sports Med 1998 Apr; 17(2):245–59. (Review of use of various substances by adolescent and young adult athletes.)

Kim S, Crutchfield C, Williams C, et al. Toward a new paradigm in substance abuse and other problem behavior prevention for youth: Youth development and empowerment approach. J Drug Educ 1998; 28(1):1–17. (Review of three decades of prevention strategies and discussion of youth development/empowerment.)

Liepman MR, Keller DM, Botelho RJ, et al. Understanding and preventing substance abuse by adolescents. Prim Care 1998 Mar; 25(1):137–62. (Review of epidemiology, stages of use, risk factors, assessment, intervention.)

Lisnov L, Harding CG, Safer LA, et al. Adolescents' perceptions of substance abuse prevention strategies. Adolescence 1998 Summer; 33(130):301–11. (Study of 719 Chicago students in grades 6–9.)

Martinez TE, Gleghorn A, Marx R, et al. Psychosocial histories, social environment, and HIV risk behaviors of injection and noninjection drug using homeless youth. J Psychoactive Drugs 1998 Jan–Mar; 30(1):1–10. (Survey of 186 homeless, runaway, and street youth ages 12–23 years.)

Miles DR, Stallings MC, Young SE, et al. A family history and direct interview study of the familial aggregation of substance abuse: The Adolescent Substance Abuse Study. Drug Alc Depend 1998 Jan; 49(2):105–14. (Study of 100 matched treatment and control probands ages 13–19 years and their family members.)

Myers MG, Stewart DG, Brown SA. Progression from conduct disorder to antisocial personality disorder following treatment for adolescent substance abuse. Am J Psychiatry 1998 Apr; 155(4):479–85. (Four-year longitudinal study of 137 patients mean age 15 years.)

Ross C, Richard L, Potvin L. One year outcome evaluation of an alcohol and drug abuse prevention program in a Quebec high school. Can J Public Health 1998 May–Jun; 89(3):166–70. (Study of 145 experimental and 179 control students in grade 7.)

Tweed SH. Intervening in adolescent substance abuse. Nurs Clin North Am 1998 May; 33(1):29–45. (Review of current patterns of use, risk factors, prevention, identification, interventions.)

Weinberg NZ, Rahdert E, Colliver JD, et al. Adolescent substance abuse: A review of the past 10 years. J Am Acad Child Adolesc Psychiatry 1998 Mar; 37(3):252–61. (Review of natural history, epidemiology, etiologic factors, co-morbidities, assessment, treatment, prevention.)

Zapata JT, Katims DS, Yin Z. A two-year study of patterns and predictors of substance use among Mexican-American youth. Adolescence 1998 Summer; 33(130):391–403. (Survey of 1805 students in grades 5–7.)

Alcohol

Duncan SC, Alpert A, Duncan TE, et al. Adolescent alcohol use development and young adult outcomes. Drug Alc Depend 1997 Dec; 49(1):39–48. (Longitudinal study of 480 youth ages 11–15 years assessed annually for four years and at ages 19–26 years.)

Kelly KJ, Edwards RW. Image advertisements for alcohol products: Is their appeal associated with adolescents' intention to consume alcohol? Adolescence 1998 Spring; 33(129):47–59. (Survey of 1058 students in grades 7, 9, or 11.)

Leichliter JS, Meilman PW, Presley CA, et al. Alcohol use and related consequences among students with varying levels of involvement in college athletics. J Am College Health 1998 Mar; 46(5):257–62. (Analysis of data from national Core Alcohol and Drug Surveys for 51,483 students.)

Ma GX, Toubbeh J, Cline J, et al. Native American adolescents' views of fetal alcohol syndrome prevention in schools. J School Health 1998 Apr; 68(4):131–6. (Survey of 163 students in grades 6–8.)

McLennan JD, Shaw E, Shema SJ, et al. Adolescents' insight in heavy drinking. J Adolesc Health 1998 May; 22(5):409–16. (Analysis of data from Arkansas Youth Risk Behavior Survey for 3395 students in grades 7–9.)

Pechansky F. Patterns of alcohol use among adolescents living in Porto Alegre, Brazil. J Psychoactive Drugs 1998 Jan–Mar; 30(1):45–51. (Survey of 950 youth ages 10–18 years.)

Scheier LM, Botvin GJ. Relations of social skills, personal competence, and adolescent alcohol use: A developmental exploratory study. J Early Adolesc 1998 Feb; 18(1):77–114. (Study of 2696 students followed from grade 6 to grade 8.)

Slater MD, Rouner D, Domenech-Rodriguez M, et al. How adolescents counterargue television beer advertisements: Implications for education efforts. J Health Educ 1998 Mar/Apr; 29(2):100–5. (Survey of 401 Anglo and Latino students responding to 4 beer and 2 non-beer ads embedded in sports and entertainment programs.)

Smeaton GL, Josiam BM, Dietrich UC. College students' binge drinking at a beach-front destination during spring break. J Am College Health 1998 Mar; 46(5):247–54. (Survey of 442 female and 341 male undergraduates.)

Williams C, Epstein JA, Botvin GJ, et al. Psychosocial determinants of alcohol use among minority youth living in public housing developments. J Dev Behav Pediatr 1998 Jun; 19(3):145–54. (Study of 624 African American and Hispanic 7th grade students.)

Marijuana

Bachman JG, Johnston LD, O'Malley PM. Explaining recent increases in students' marijuana use: Impacts of perceived risks and disapproval, 1976 through 1996. Am J Public Health 1998 Jun; 88(6):887–92. (Analysis of national data from Monitoring

the Future project for high school senior classes from 1976 to 1996 and 8th and 10th grade classes from 1991 to 1996.)

Crowley TJ, MacDonald MJ, Whitmore EA, et al. Cannabis dependence, withdrawal, and reinforcing effects among adolescents with conduct symptoms and substance use disorders. Drug Alc Depend 1998 Mar; 50(1):27–37. (Study of 229 patients ages 13–19 years.)

Ewing BT. High school athletes and marijuana use. J Drug Educ 1998; 28(2):147–57. (Analysis of data from National Longitudinal Survey of Youth for 1458 students in grades 9–12.)

Rosenbaum M. "Just say know" to teenagers and marijuana. J Psychoactive Drugs 1998 Apr–Jun; 30(2):197–203. (Commentary on marijuana use and current drug education.)

Van Etten ML, Neumarky D, Anthony JC. Initial opportunity to use marijuana and the transition to first use: United States, 1979–1994. Drug Alc Depend 1997 Dec; 49(1):1–7. (Analysis of data from U.S. National Household Surveys on Drug Abuse.)

Tobacco

Ammerman SD. Helping kids kick butts. Contemp Pediatr 1998 Feb; 15(2):64–77. (Review.)

Botvin GJ, Epstein JA, Botvin EM. Adolescent cigarette smoking: Prevalence, causes, and intervention approaches. AM:STARS 1998 Jun; 9(2):299–313. (Review of epidemiology, etiologic factors, prevention strategies, practical considerations.)

Cummings KM, Hyland A, Saunders-Martin T, et al. Evaluation of an enforcement program to reduce tobacco sales to minors. Am J Public Health 1998 Jun; 88(6):932–6. (Study of sales to minors in 319 outlets in 6 pairs of New York communities.)

Distefan JM, Gilpin EA, Choi WS, et al. Parental influences predict adolescent smoking in the United States, 1989–1993. J Adolesc Health 1998 Jun; 22(6):466–74. (Analysis of data from Teenage Attitudes and Practices Surveys.)

Dovell RA, Mowat DL, Dorland J, et al. Tobacco access to youth: Beliefs and attitudes of retailers. Can J Public Health 1998 Jan–Feb; 89(1):17–21. (Phone survey of 151 Ontario retail outlets.)

Everett SA, Husten CG, Warren CW, et al. Trends in tobacco use among high school students in the United States, 1991–1995. J School Health 1998 Apr; 68(4):137–40. (Analysis of data from 1991, 1993, and 1995 Youth Risk Behavior Surveys for 12,227, 16,296, and 10,904 students in grades 9–12.)

Flint AJ, Novotny TE. Trends in black/white differences in current smoking among 18- to 24-year-olds in the United States, 1983–1993. Am J Prev Med 1998 Jan; 14(1):19–24. (Analysis of data from National Health Interview Surveys for 33,838 individuals.)

Gemson DH, Moats HL, Watkins BX, et al. Laying down the law: Reducing illegal tobacco sales to minors in central Harlem. Am J Public Health 1998 Jun; 88(6):936–9. (Study of sales to 12–13-year-olds at 152 stores.)

Griesler PC, Kandel DB, Davies M. Maternal smoking in pregnancy, child behavior problems, and adolescent smoking. J Res Adolesc 1998; 8(2):159–85. (Longitudinal study of 187 mothers ages 34–35 years with first-born children ages 9–17 years.)

Hu T, Lin Z, Keeler TE. Teenage smoking, attempts to quit, and school performance. Am J Public Health 1998 Jun; 88(6):940–3. (Analysis of data from 1990 California Youth Tobacco Survey for 5028 youth ages 12–17 years.)

King C, Siegel M, Celebucki C, et al. Adolescent exposure to cigarette advertising in magazines: An evaluation of brand-specific advertising in relation to youth readership. JAMA 1998 Feb; 279(7):516–20. (Study of readers age 12–17 and age 18–24 years of 39 popular American magazines.)

Lamkin L, Houston TP. Nicotine dependency and adolescents: Preventing and treating. Prim Care 1998 Mar; 25(1):123–35. (Review of trends in tobacco use, research on cessation, interventions.)

McCormick L, Tompkins NO. Diffusion of CDC's guidelines to prevent tobacco use and addiction. J School Health 1998 Feb; 68(2):43–5. (Survey of state and local education agencies in 15 states.)

Office on Smoking and Health, Centers for Disease Control and Prevention. Tobacco use among high school students—United States, 1997. J School Health 1998 May; 68(5):202–4. (Analysis of data from 1997 Youth Risk Behavior Survey for 16,262 students in grades 9–12.)

Pierce JP, Choi WS, Gilpin EA, et al. Tobacco industry promotion of cigarettes and adolescent smoking. JAMA 1998 Feb; 279(7): 511–5. (Telephone survey of 1752 California adolescents in 1993 and again in 1996.)

Rojas NL, Killen JD, Haydel KF, et al. Nicotine dependence among adolescent smokers. Arch Pediatr Adolesc Med 1998 Feb; 152(2):151–6. (Survey of 2197 California 10th grade students.)

Sargent JD, Mott LA, Stevens M. Predictors of smoking cessation in adolescents. Arch Pediatr Adolesc Med 1998 Apr; 152(4): 388–93. (Study of 1384 New Hampshire high school students ages 12–18 years assessed yearly for 3 years.)

Tomar SL, Giovino GA. Incidence and predictors of smokeless tobacco use among US youth. Am J Public Health 1998 Jan; 88(1):20–6. (Analysis of data from 1989 Teenage Attitudes and Practices Survey and its 1993 follow-up for 7960 youth ages 12–18 years.)

Vartiainen E, Paavola M, McAlister A, et al. Fifteen-year follow-up of smoking prevention effects in the North Karelia Youth Project. Am J Public Health 1998 Jan; 88(1):81–5. (Study of Finns at age 13, 15, 16, 17, 21, and 28 years.)

Wang MQ, Fitzhugh EC, Eddy JM, et al. Tobacco use among school adolescents: National sociodemographic risk profiles. J Health Educ 1998 May/Jun; 29(3):174–8. (Analysis of data from 1993 Teenage Attitudes and Practices Survey for 8725 students ages 12–18 years.)

SURGERY/TRAUMA

American Academy of Pediatrics Committee on Injury and Poison Prevention and Committee on Sports Medicine and Fitness. In-line skating injuries in children and adolescents. Pediatrics 1998 Apr; 101(4):720–2. (Position statement.)

Dunn KA, Runyan CW, Cohen L, et al. Teens at work: A statewide study of jobs, hazards, and injuries. J Adolesc Health 1998 Jan; 22(1):19–25. (North Carolina survey of 562 youth ages 14–17 years.)

Marven SS, Smith CM, Claxton D, et al. Pulmonary function, exercise performance, and growth in survivors of congenital diaphragmatic hernia. Arch Dis Child 1998 Feb; 78(2):137–42. (Study of 19 survivors and 19 matched controls ages 7–16 years.)

Miller TR, Spicer RS. How safe are our schools? Am J Public Health 1998 Mar; 88(3):413–8. (Analysis of data from National Health Interview Surveys from 1987 through 1992 for youth ages 5–10, 11–14, and 15–19 years.)

Tesler MD, Holzemer WL, Savedra MC. Pain behaviors: Postsurgical responses of children and adolescents. J Pediatr Nurs 1998 Feb; 13(1):41–7. (Study of 37 patients ages 8–17 years.)